

CONTENTS

Biotization of <i>Pseudomonas fluorescens</i> isolates in groundnut (<i>Arachis hypogaea</i> L.) against root-knot nematode, <i>Meloidogyne arenaria</i> --P. Kalaiarasan, P.L. Lakshmana and R. Samiyappan	--	1
Population dynamics of plant parasitic nematodes in different cropping sequences under wetland system --P. Vetrivelkalai and S. Subramanian	--	6
Plant parasitic nematode populations as influenced by onion based cropping sequences --P. Vetrivelkalai and S. Subramanian	--	10
Differential induction of chitinase activity in resistant and susceptible sugarcane clones infected by <i>Pratylenchus zeae</i> --T. Kathiresan and Usha K. Mehta	--	15
Biocontrol efficacy of entomopathogenic nematodes on <i>Spodoptera litura</i> (Lepidoptera: Noctuidae) in blackgram --R. Umamaheswari, M. Sivakumar and S. Subramanian	--	19
Studies on the method of application of the fungus, <i>Paecilomyces lilacinus</i> against root-knot nematode on tomato --M. Shanmuga Priya and S. Kumar	--	23
Dose optimization of <i>Paecilomyces lilacinus</i> for the control of <i>Meloidogyne incognita</i> on tomato --M. Shanmuga Priya and S. Kumar	--	27
Two new species of mononch (Nematoda: Mononchida) from soil around the rhizosphere of bamboo from Manipur --Ch. Dhanachand, H. Romoni and M. Pramodini	--	32
Efficacy of organic amendment in management of root-knot nematode <i>Meloidogyne incognita</i> on okra --B.B. Bhosle, Mukesh Sehgal, S.N. Puri, H.R. Sardana and D.K. Singh	--	37
Cellular response of blackgram roots to combined infection of <i>Meloidogyne incognita</i> and <i>Fusarium oxysporum</i> --S.N. Mahapatra and P.K. Swain	--	41
Management of foliar nematode, <i>Aphelenchoides besseyi</i> infecting tuberose in West Bengal, India --M.R. Khan, S. Shit, A.K. Pal and B. Biswas	--	44
Species-specific enzyme phenotypes of <i>Steinernema carpocapsae</i> , <i>S. glaseri</i> and <i>S. thermophilum</i> (Nematoda: Steinernematidae) --Sudershan Ganguly, Jola Pandey and Kabindra Singh Rathour	--	48
Preliminary evaluation of the management of <i>Meloidogyne incognita</i> using biodegradable and synthetic matrices for carbofuran formulations --M.A. Radwan, H.S. Ibrahim, S.M.I. Kassem and M.M. Abu-Elamayem	--	54
Nutrient uptake as affected by inorganic fertilizers under <i>Meloidogyne incognita</i> infested conditions in cotton --K.K. Verma and R.K. Jain	--	60
Effect of different bioinoculants, organic amendments and pesticides on the management of <i>Meloidogyne incognita</i> - <i>Fusarium solani</i> disease complex on tomato cv. K-25 --Akhtar Haseeb, Vipin Kumar, Prabhat Kumar Shukla and Abrar Ahmad	--	65
A new species <i>Discocriconemella waitha</i> from Manipur with redescription and male report of <i>Discocriconemella spermata</i> Mohilal and Dhanachand, 1998 --M. Pramodini, N. Mohilal and Ch. Dhanachand	--	70
Evaluation of salicylic acid as a systemic resistance inducer against <i>Meloidogyne incognita</i> on tomato cv. Co3 --J. Jayakumar, G. Rajendran and S. Ramakrishnan	--	77
Effect of crop sequences on incidence of root-knot and wilt disease complex of jute caused by <i>Meloidogyne incognita</i> and <i>Ralstonia solanacearum</i> --K. Hazarika, F. Rahman and L.C. Bora	--	81
Out break of root-knot nematode (<i>Meloidogyne graminicola</i>) disease in rice and farmers' perceptions --J.S. Prasad, Vishakanta and Gubbaiah	--	85
Community structure of plant parasitic nematodes in Champawat district of Uttaranchal, India -- Kabindra Singh Rathour, Jola Pandey and Sudershan Ganguly	--	89
Mass production and viability of an egg parasitic fungus, <i>Paecilomyces lilacinus</i> --S.C. Dhawan, Sumeet, M.N. Tripathi and N. Prakash Babu	--	94

(continued on inner cover)

Community analysis of plant parasitic nematodes associated with pulse crops in Hamirpur district (Uttar Pradesh) --S.S. Ali, Rashid Pervez, Azra Shaheen and R. Ahmad	--	99
Occurrence and distribution on reniform nematode, <i>Rotylenchulus reniformis</i> in Udaipur (Rajasthan) India --G.M. Lone, A.U. Siddiqui, A. Parihar, S.S. Sharma and S.D. Masoodi	--	103
Nematicidal activity of essential oils and their components against the root-knot nematode <i>Meloidogyne incognita</i> --Archana Sinha, R.C. Maheshwari, Prem Dureja and Vijayalakshmi Mojumder	--	109
Molecular characterization of <i>Steinernema thermophilum</i> based on RFLP and sequence of ITS region of rDNA --Sudershan Ganguly and Jola Pandey	--	115
SHORT COMMUNICATIONS		
Preliminary screening of benzalaniline and its C-phenyl derivatives for nematicidal activity --Balbir Kaur Matharu, V.K. Kaul and M.R. Manrao	--	122
Pathogenicity of <i>Meloidogyne incognita</i> on <i>Coleus forskohlii</i> Briq. --M. Senthamarai, K. Poornima and S. Subramanian	--	123
Management of root-knot nematode, <i>Meloidogyne javanica</i> infecting tomato --V.K. Singh	--	126
Management of root-knot nematode, <i>Meloidogyne incognita</i> infecting cauliflower --V.K. Singh	--	127
Extracting efficiency of different versions of Baermann's Funnel Technique --K.R.Dabur and A.S. Taya	--	128
Survival, infectivity and reproduction of entomopathogenic nematodes in spray adjuvants --P. Priya and S. Subramanian	--	129
Nematophagus fungus associated with pine trees in Manipur --L. Victoria, N. Mohilal and Ch. Dhanachand	--	130
Phytotherapeutic management of root-knot nematode (<i>Meloidogyne incognita</i>) through peels of <i>Citrus reticulata</i> (Orange) --Ranjana Saxena and Shalini Gupta	--	131
Effect of culture filtrates of <i>Pseudomonas fluorescens</i> and <i>Trichoderma viride</i> and root exudates of <i>Glomus mosseae</i> on hatching of <i>Rotylenchulus reniformis</i> --S. Subramanian and K. Senthamizh	--	133
Bio-management of <i>Meloidogyne incognita</i> on coleus, <i>Solenostemon rotundifolius</i> by integrating solarization, <i>Paecilomyces lilacinus</i> , <i>Bacillus macerans</i> and neemcake --M.S. Nisha and M.S. Sheela	--	136
Effect of culture filtrates of <i>Pseudomonas fluorescens</i> and <i>Trichoderma viride</i> and root exudates of <i>Glomus mosseae</i> on mortality of <i>Rotylenchulus reniformis</i> --S. Subramanian and K. Senthamizh	--	138
Management of reniform nematode, <i>Rotylenchulus reniformis</i> on cowpea (<i>Vigna unguiculata</i> L.) with plant products --Budh Ram and B.L. Baheti	--	140
<i>In vitro</i> mass production technology for the entomopathogenic nematode, <i>Steinernema glaseri</i> --S. Prabhu, G. Rajendran and S. Subramanian	--	142
Effect of leaf extracts of <i>Acalypha indica</i> , <i>Cassia fistula</i> and <i>Solanum torvum</i> on <i>Pratylenchus coffeae</i> --P. Sundraraju and V. Saritha	--	144
Pathogenicity of <i>Paraphelenchus</i> species of milky mushroom (<i>Calocybe indica</i>) and its multiplication on other fungi --Rajesh Vats, Surjit Singh and Harish K. Bajaj	--	146
Effect of nematicides and organic amendments in the management of <i>Meloidogyne incognita</i> in nursery tea --S. Kalita and B.C. Bora	--	148
Evaluation of wheat and barley germplasm against the cereal cyst nematode, <i>Heterodera avenae</i> --Pankaj, B.D. Yadav, Anil Sirohi, Veer Kishor, Deepika Rohatgi, H.S. Gaur and R.J. Meena	--	150
Efficacy of oil cakes as seed dressing treatment for the management of root-knot nematode, <i>Meloidogyne incognita</i> affecting chickpea --Y.S. Yadav, A.U. Siddiqui and Aruna Parihar	--	151
Effect of oil cakes as soil application in management of root-knot nematode, <i>Meloidogyne incognita</i> affecting chickpea --Y.S. Yadav, A.U. Siddiqui and Aruna Parihar	--	153
Nematode communities associated with litchi at Muzaffarnagar, Uttar Pradesh, India --M. Saha, M. Lal and M. Singh	--	154