

CONTENTS

Recent taxonomic status of entomopathogenic nematodes: a review --Sudershan Ganguly	-- 158
Pathogenic potential of <i>Meloidogyne incognita</i> on <i>Mentha arvensis</i> cv. Gomti --K. Perveen, A. Haseeb and P.K. Shukla	-- 177
Nematode-fungal disease complex involving <i>Meloidogyne incognita</i> and <i>Macrophomina phaseolina</i> on <i>Coleus forskohlii</i> Briq. --M. Senthamarai, Poornima and S. Subramanian	-- 181
Effect of cropping sequences on the population dynamics of plant parasitic nematodes under dryland systems --P. Vetrivelkai and S. Subramanian	-- 185
Effect of <i>Glomus fasciculatum</i> on nutritional and biochemical changes in brinjal infested by <i>Meloidogyne incognita</i> --Aparajita Borah and P.N. Phukan	-- 189
Control of plant parasitic nematodes associated with linseed by oil seed cakes and ploughing --Suhail Anver	-- 192
Effect of vermiwash on nematodes prevalent in coconut based high-density multispecies cropping system --J. Gulsar Banu and Rohini Iyer	-- 195
Chitin and chitinolytic biocontrol agents for the management of root knot nematode, <i>Meloidogyne arenaria</i> in groundnut (<i>Arachis hypogaea</i> L.) cv. Co3 --P. Kalaiarasan, PL. Lakshmanan, G. Rajendran and R. Samiyappam	-- 200
Bio-management of root-knot nematode, <i>Meloidogyne incognita</i> on <i>Coleus forskohlii</i> Briq. --M. Senthamarai, K. Poornima and S. Subramanian	-- 206
Effect of graded doses of nitrogen and VAM on Herbage, oil yield and nematode population on Patchouli (<i>Pogostemon patchouli</i> Pellet) --M. Sumathi, S. Parthiban and Stephen D. Samuel	-- 209
Efficiency of some native isolates of <i>Saccharomyces cerevisiae</i> , <i>Trichoderma harzianum</i> and <i>T. reesei</i> as bioagents and vdate against <i>Helicotylenchus exallus</i> and <i>Pratylenchus zeae</i> infesting jasmine in Egypt --A.E. Ismail, S.A. Hasabo and W.M.A. El-Nagdi	-- 214
Fluctuation of nematode populations associated with banana plantation in Medinipur district, West Bengal, India --V.V. Gantait, T. Bhattacharya and A. Chatterjee	-- 223
Community structure of plant parasitic nematodes in banana plantations of Andhra Pradesh, India --P. Sundararaju	-- 226
Biomangement of nematodes infesting banana by bacterial endophytes (<i>Bacillus subtilis</i>) --E.I. Jonathan and R. Umamaheshwari	-- 230
Bioefficacy of native plant growth promoting rhizobacteria against <i>Meloidogyne incognita</i> and <i>Phytophthora capsici</i> disease complex in betelvine --E.I. Jonathan, R. Umamaheshwari and P. Bommaraju	-- 234
Isolation of native <i>Xenorhabdus</i> bacteria from <i>Steinernema</i> spp. and role of their exo and endo metabolites for suppression of root-knot nematodes (<i>Meloidogyne</i> spp.) on tomato --R.V. Vyas, A.B. Maghodia, Biren A. Patel and D.J. Patel	-- 241
Morphometric variations in Indian populations of <i>Heterodera cajani</i> Koshy, 1967. I-second stage juveniles and males --M. Abdollahi, A.K. Ganguly and K.K. Kaushal	-- 247
Biochemical changes in banana plants induced by <i>Pratylenchus coffeae</i> and <i>Meloidogyne incognita</i> --P. Sundararaju and K. Pandisuba	-- 256
Pathogenicity and evaluation of plant root extracts on root-knot nematode infecting tomato --T.I. Olabiyi	-- 260
Role of <i>Trichoderma harzianum</i> and neem cake separately and in combination against root-knot nematode on tomato --Sunil Kumar and Anju S. Khanna	-- 264
Sources of nematode inoculum in white button and milky mushroom --Anju S. Khanna, R. Chandran A. and Sunil Kumar	-- 267
SHORT COMMUNICATIONS	
Influence of previous crop on the density and frequency of plant parasitic nematodes associated with groundnut --K. Senthamizh, M. Sivakumar and C.V. Sivakumar	-- 270
<i>Gracilacus vitecus</i> sp.n. and record of <i>G. raskii</i> Phukan & Sanwal from Manipur, India --M. Pramodini, N. Mohilal and Ch. Dhanachand	-- 272
Nematodes of cut flowers in Tamil Nadu, India --Nirmal Johnson, S.B.J. Jaya Kumar, G. Rajendran and S. Ramakrishanan	-- 276

Pathogenic potential of root-knot nematode, <i>Meloidogyne arenaria</i> on groundnut (<i>Arachis hypogaea</i> L.) --P. Kalaierasan, G. Rajendran and PL. Lakshmanan	-- 278
Phylogenetic relationships of <i>Steinernema thermophilum</i> with other species of the genus based on ITS region of rDNA --Sudershan Ganguly, Jola Pandey, K.S. Rathour and V.G.Malathi	-- 280
Nematicidal activity of thio substituted ketimines --Monika Jolly, V.K. Kaul and M.R. Manrao	-- 281
Current status of entomopathogenic nematodes occurring in India along with new record of <i>Steinernema feltiae</i> (Filipjev) Wouts <i>et al.</i> from Kerala --Sudershan Ganguly, K.S. Rathour, Jola Pandey and V.K.Sosamma	-- 283
Virulence of <i>Heterorhabditis indica</i> to the grubs of lucerne weevil, <i>Hypera postica</i> (Coleoptera; Curculionidae) --N.K. Shah and M.I. Azmi	-- 285
Effect of various levels of fertilizer in the interaction of <i>Glomus fasciculatum</i> and <i>Meloidogyne incognita</i> on greengram --Aparajita Borah, P.P. Neog and A.K. Sinha	-- 287
Effect of different levels of moisture on the interaction of <i>Glomus fasciculatum</i> with <i>Meloidogyne inognita</i> on blackgram --Aparajita Borah, P.P. Neog and A.K. Sinha	-- 288
Effect of <i>Vicia faba</i> on the morphometrics of root-knot nematode, <i>Meloidogyne incognita</i> --S.P. Singh and Veena Singh	-- 290
Reaction of pulse genotypes for resistance to pigeonpea cyst nematode, <i>Heterodera cajani</i> Koshy, 1967 --P. Kalaierasan, M. Sivakumar and S. Subramanian	-- 291
Screening of okra varieties for resistance against <i>Meloidogyne incognita</i> --M.S. Sheela, Jiji, Malu Ravi and Shaiju Simon	-- 292
Screening of vegetable crop varieties for resistance to root-knot nematode (<i>Meloidogyne incognita</i>) --P. Harinath Naidu, V. Haritha and M. John Sudheer	-- 293
Duranta - a new host for root-knot nematode (<i>Meloidogyne incognita</i>) --P. Harinath Naidu and V. Haritha	-- 294
Comparative efficacy of nematicides and neem cake for the management of <i>Meloidogyne incognita</i> in french bean --A. Hussain and B.C. Bora	-- 295
Assessment of avoidable yield loss on <i>Coleus forskohlii</i> due to <i>Meloidogyne incognita</i> --M. Senthamarai, K. Poornima and S. Subramanian	-- 296
Effect of host plant on mass production of <i>Pasteuria penetrans</i> --C. Gomathi, S. Kumar and S. Subramanian	-- 297
Efficacy of carbofuran, bavistin, neem, <i>Trichoderma harzianum</i> and <i>Aspergillus niger</i> against <i>Meloidogyne incognita</i> and <i>Fusarium oxysporum</i> f. sp. <i>vasinfectum</i> disease complex on okra --Syed Abuzar and Akhtar Haseeb	-- 299
Field performance of cereal cyst nematode resistant wheat variety raj MR1 in Haryana --R.S. Kanwar and S.N. Nandal	-- 301
Reaction of <i>Heterodera avenae</i> resistant varieties of wheat and barley against rice root-knot nematode, <i>Meloidogyne graminicola</i> --R.S. Kanwar, K.R. Dabur and S.N. Nandal	-- 302
Searching resistance and tolerance in wheat cultivars against cereal cyst nematode, <i>Heterodera avenae</i> Woll. --Brigbhan and R.S. Kanwar	-- 303
Influence of <i>Pseudomonas fluorescens</i> on histopathological changes caused by <i>Meloidogyne arenaria</i> in groundnut (<i>Arachis hypogaea</i> L.) --P. Kalaierasan, PL. Lakshmanan and R. Samiyappam	-- 305
Integrated management of root-knot nematode in tomato nursery through botanical plant materials --S.K. Patel, H.V. Patel and A.D. Patel	-- 307
Effect of some compatible biocontrol agents along with mustard cake and furadon on <i>Meloidogyne incognita</i> infecting tomato plant --Pranod Kumar Tripathi and C.S. Singh	-- 309
Record of the rice cyst nematode, <i>Heterodera oryzae</i> in Kerala, India --V.K. Sosamma and P.K. Koshy	-- 311
Infestation of <i>Radopholus similis</i> with <i>Vorticella</i> from Kerala --V.K. Sosamma	-- 312
Efficacy of dry powder of <i>Adhatoda vasica</i> as soil amendment against root-knot nematode on brinjal --L. Joymati Devi	-- 312
<i>Pronycolaimus shamimi</i> gen. n., sp. n. (Dorylaimida: Nematoda) from India --Vinita Sharma and Qaiser H. Baqri	-- 314
Book review	-- 317