

## CONTENTS

Comparative effects of culture filtrate of soil-borne fungi on mortality and infectivity of juveniles of <i>Meloidogyne incognita</i> --M.A. Radwan	-- 109
Jatropha: An efficient organic amendment for the management of root-knot nematode, <i>Meloidogyne incognita</i> --P. Kalaiarasan, M. Senthamarai, D. Ramesh and M. John Sudheer	-- 115
Effect of fly ash amended soil on the development of root-knot nematode, <i>Meloidogyne incognita</i> and growth of ivy gourd, <i>Coccinea cordifolia</i> --Tanweer Azam, Hisamuddin, T. Niyaz and M.I. Robab	-- 119
Occurrence of entomopathogenic nematodes in varied agro-climatic zones of Rajasthan --A.K. Maru, A.U. Siddiqui, A. Parihar and S.K. Sharma	-- 123
Effect of opportunistic fungi on eggmass parasitization and mortality of juveniles of <i>Meloidogyne incognita</i> <i>In vitro</i> --M. Reddi Kumar, J. Sailaja Rani and M. John Sudheer	-- 127
Role of phenolics and enzymes in imparting resistance to rice plants against root-knot nematode, <i>Meloidogyne graminicola</i> --C.D. Mishra and K.C. Mohanty	-- 131
Management of root-knot nematode, <i>Meloidogyne incognita</i> by <i>Paecilomyces lilacinus</i> -influence of soil moisture and soil temperature --R. Krishnamoorthi and S. Kumar	-- 135
Mechanism of resistance in banana cultivars against root lesion nematode, <i>Pratylenchus coffeae</i> --A. Nithya Devi, V. Ponnuswami, P. Sundararaju, K. Soorianathasundaram, S. Sathiamoorthy, S. Uma and I. Van den Bergh	-- 138
Bioefficacy of entomopathogenic nematodes, <i>Steinernema carpocapsae</i> and <i>Heterorhabditis indica</i> against third instar larvae of <i>Spodoptera litura</i> --S. Raveendranath, P.V. Krishnayya, P. Arjuna Rao, K.V.M. Krishna Murthy and S.S. Hussaini	-- 145
Phenylalanine ammonia lyase and total phenol content in resistant banana to <i>Pratylenchus coffeae</i> --A. Nithya Devi, V. Ponnuswami, P. Sundararaju, K. Soorianathasundaram, S. Sathiamoorthy, S. Uma and I. Van den Bergh	-- 149
Histopathology of cowpea roots co-infected with root-knot nematode, <i>Meloidogyne incognita</i> and wilt fungus, <i>Fusarium oxysporum</i> --Satyandra Singh, S.C. Dhawan and B.K. Goswami	-- 156
Nemic population dynamics in hybrid tomato, sweet pepper and hybrid cucumber under polyhouse cultivation --H.K. Sharma, Pankaj, H.S. Gaur and Balraj Singh	-- 161
Life cycle, varietal reaction, biochemical alteration and histopathology of rice root-knot nematode, <i>Meloidogyne graminicola</i> --P. Senthilkumar, S. Ramakrishnan and E.I. Jonathan	-- 165
Role of reniform nematode, <i>Rotylenchulus reniformis</i> in reducing disease-complex on tomato caused by root-knot nematode, <i>Meloidogyne incognita</i> and wilt fungus, <i>Fusarium oxysporum</i> f.sp. <i>lycopersici</i> --Satyandra Singh, B.K. Goswami and H.K. Sharma	-- 172
Evaluation of essential oil products of medicinal plants on egg hatching and larval mortality of <i>Meloidogyne incognita</i> --L. Joymati Devi	-- 176

Efficacy of organic amendments in combination with the nematicide for management of <i>Meloidogyne incognita</i> in brinjal --J. Saikia, Aparajita Borah and B.C. Bora	--	179
Random amplified polymorphic DNA markers for prediction of resistance in tomato against root-knot nematode, <i>Meloidogyne incognita</i> --Anil Sirohi, Atul Kakrana, Pankaj, Deepika Rohatgi and R.K. Jain	--	182
Influence of 28-homobrassinolide on development of second-stage juveniles of <i>Meloidogyne incognita</i> (Kofoid & White) Chitwood --P. Ohri, S.K. Sohal, R. Bhardwaj and U.R. Khurma	--	188
Combined effect of inorganic supplements with <i>Trichoderma viride</i> and <i>Kalisena</i> against root-knot nematode infecting okra --H.K. Sharma, Pankaj, Jagan Lal and Satyandra Singh	--	192
<b>SHORT COMMUNICATIONS</b>		
Role of plant parts and their extracts in the management of <i>Meloidogyne incognita</i> infecting pigeonpea --K.D. Upadhyay, Kusum Dwivedi and S.K. Uttam	--	197
Plant-parasitic nematodes associated with medicinal coleus ( <i>Coleus forskohlii</i> ) in Tamil Nadu, India --N. Seenivasan and K. Devrajan	--	199
Report of root-knot nematode, <i>Meloidogyne incognita</i> on pomegranate, <i>Punica granatum</i> L. from Andhra Pradesh --M. John Sudheer, P. Kalaiarasan and M. Senthamarai	--	201
Effect on viability and infectivity of entomopathogenic nematodes (Meghalaya isolates) --Gitanjali Devi	--	202
Nematicidal activity of bis-ketimines against <i>Ditylenchus myceliophagus</i> --M.R. Manrao, Gurbir Kaur and V.K. Kaul	--	205
Community analysis of plant parasitic nematodes in Chittoor district of Andhra Pradesh --P. Harinath Naidu, V. Haritha and M. John Sudheer	--	207
Status of wheat seed gall nematode, <i>Anguina tritici</i> in Haryana --R.S. Kanwar and S.N. Nandal	--	212
Preliminary evaluation of the entomopathogenic nematodes on the root grub, <i>Basilepta fulvicorne</i> in cardamom --I. Cannayane, J. Gulsar Banu, S. Subramanian and D.S. Rajavel	--	213
Occurrence of <i>Steinernema masoodi</i> in Aligarh and its pathogenicity against six economically important insect pests-- --Mujeebur Rahman Khan, Uzma Khan and Tariq H. Askary	--	215
Effect of aqueous extracts of plant products on hatching of <i>Meloidogyne incognita</i> Race 2 infecting tomato --P. Harinath Naidu, V. Haritha and M. John Sudheer	--	217
Estimation of losses due to plant parasitic nematodes on different crops in India --R.K. Jain, K.N. Mathur and R.V. Singh	--	219
Community analysis of plant parasitic nematodes associated with vegetable crops in selected districts of Andhra Pradesh --G.M.V. Prasada Rao, M. John Sudheer and P. Priya	--	221
Reaction of chickpea germplasm against root-knot nematode, <i>Meloidogyne incognita</i> --S.M. Yadav, Mukesh Sehgal and Y.S. Yadav	--	223