

## CONTENTS

Effect of plant leaf ash on the control of <i>Meloidogyne incognita</i> in soybean ( <i>Glycine max</i> (L.) Merrill.) --C.C. Ononuju and U.C. Kpadobi	-- 1
Management of root-knot nematode, <i>Meloidogyne incognita</i> using bio-control agents on medicinal coleus, <i>Coleus forskohlii</i> Briq. --M. Senthamarai, K. Poornima and S. Subramanian	-- 5
Relative susceptibility of cultivated crops to reniform nematode, <i>Rotylenchulus reniformis</i> and its host range in and around Udaipur (Rajasthan), India --G.M. Lone, A.U. Siddiqui, M.I.S. Walliullah, Aruna Parihar and S.S. Sharma	-- 9
Management of <i>Globodera rostochiensis</i> and <i>G. pallida</i> in potato by antagonistic crop, mustard in Nilgiris --K. Devrajan and P. Balasubramanian	-- 15
Influence of <i>Pseudomonas fluorescens</i> on histopathological changes caused by <i>Meloidogyne arenaria</i> in groundnut ( <i>Arachis hypogaea</i> L.) --P. Kalairasan, P.L. Lakshmanan, R. Samiyappan and M. John Sudheer	-- 18
Population dynamics of root-knot nematode under different cropping sequences at farmers fields --I.J. Paruthi	-- 21
Management approaches for tuberose foliar nematode, <i>Aphelenchoides besseyi</i> in West Bengal, India --M.R. Khan, B. Pathak, S. Ghosh and K. Ghoshdostidar	-- 25
Nematode-Fungal disease complex involving <i>Meloidogyne incognita</i> and <i>Macrophomina phaseolina</i> on medicinal coleus, <i>Coleus forskohlii</i> Briq. --M. Senthamarai, K. Poornima, S. Subramanian and M. John Sudheer	-- 30
Engineering genetic resistance against root-knot nematode, <i>Meloidogyne incognita</i> in tomato using a antifungal rice chitinase gene ( <i>chi 11</i> ) --P. Kalairasan, M. Sivakumar and D. Sudhakar	-- 34
Growth and anatomy of green poppy ( <i>Papaver rhoeas</i> ) infected with <i>Meloidogyne incognita</i> --Tabassum Niyaz and Hisamuddin	-- 42
Bioefficacy of <i>Pseudomonas fluorescens</i> on burrowing nematode, <i>Radopholus similis</i> in banana --P. Senthilkumar, E.I. Jonathan and R. Samiyappan	-- 46
Community analysis of plant parasitic nematodes associated with agricultural crops in Junagadh district of Gujarat and Diu-Union territory --A.D. Patel, Bindu K. Panickar, B.A. Patel and D.J. Patel	-- 53
Field application of biocontrol agent, <i>Pseudomonas fluorescens</i> for the management of burrowing nematode, <i>Radopholus similis</i> in banana --P. Senthilkumar, E.I. Jonathan and R. Samiyappan	-- 57
Nematicidal activity of 1-phenylethylidenemalononitrile derivatives --Monika Jolly, M.R. Manrao and V.K. Kaul	-- 62
Biological control of <i>Heterodera cajani</i> in pigeonpea by <i>Trichoderma harzianum</i> and <i>Pochonia chlamydosporia</i> --S. Kumar and S. Prabhu	-- 65
Diversity and community structure of major plant parasitic nematodes in selected districts of Andhra Pradesh, India --M. John Sudheer, P. Kalairasan, M. Senthamarai, S. Prabhu, G.M.V. Prasad Rao, P. Priya, P. Harinatha Naidu, V. Haritha, M. Reddy Kumar and Sailaja Rani	-- 68

Occurrence, distribution and importance of plant parasitic nematodes associated with litchi plantations in north Tripura district --R.C. Nath, B.C. Sinha, B. Mukherjee and M.K. Dasgupta	-- 75
Molecular characterization of Indian populations of maize cyst nematode, <i>Heterodera zae</i> using PCR-RFLP of ribosomal DNA --UmaRao, Tony Grace, K.K. Kaushal and Vangapandu Sashi	-- 81
Comparative efficacy of <i>Trichoderma harzianum</i> and <i>Pseudomonas fluorescens</i> against <i>Meloidogyne incognita</i> and <i>Ralstonia solanacearum</i> complex in brinjal --Liza Barua and B.C. Bora	-- 86
Effect of culture filtrates of <i>Paecilomyces lilacinus</i> on the mortality and hatching of root-knot nematode, <i>Meloidogyne incognita</i> --M. Reddi Kumar, J. Sailaja Rani and M. John Sudheer	-- 90
Two new species of <i>Xenocriconemella</i> De Grisse and Loof, 1965 (Tylenchida: Criconematoidea) --Sudershan Ganguly, M. Singh, M. Lal and K.S. Rathour	-- 94
Root-knot nematode, <i>Meloidogyne incognita</i> and salicyclic acid induce production of PR-proteins in tomato --Anil Sirohi, Pankaj, Deepika Rohatgi and Divya Chaudhary	-- 98

## SHORT COMMUNICATIONS

Species and races of root-knot nematode prevalence in Nagaon district of Assam --Zakir Hussain and B.C. Bora	-- 104
Management of root-knot nematode, <i>Meloidogyne incognita</i> infecting brinjal through farmers participatory mode --Pankaj, H.K. Sharma and A.N. Srivastava	-- 105
Pathogenicity of <i>Pratylenchus</i> spp. on ginger and maize in the Indian Himalayas --A. Thapa, N. Gurung, Y.R. Samra, J. Bridge and G.V.H. Jackson	-- 107
Response of resistant cultivars of wheat and barley to <i>Heterodera avenae</i> --S.P. Bishnoi, S.M. Yadav and Ramesh Chand	-- 110
Management of root-knot nematode, <i>Meloidogyne incognita</i> infecting pigeonpea --V.K. Singh	-- 112
Reaction of pigeon pea genotypes against pigeon pea cyst nematode, <i>Heterodera cajani</i> --S.C. Dhawan and Satyandra Singh	-- 113
Comparative efficacy of neem leaves, bark and kernel extracts on mortality of root-knot nematode, <i>Meloidogyne incognita</i> --Krishan Singh and K.R. Dabur	-- 115
Community analysis of nematodes in village Kawas, Barmer, Rajasthan --Pukhraj Kadela	-- 117
Bio-management of root-knot nematode, <i>Meloidogyne incognita</i> by <i>Pochonia chlamydosporia</i> and <i>Pseudomonas fluorescens</i> on brinjal in farmer's field --S.C. Dhawan, Satyandra Singh and Anju Kamra	-- 119
Host status of different seed spices against root-knot nematode ( <i>Meloidogyne incognita</i> ) --R.K. Jain and I.J. Paruthi	-- 121
Level of awareness of phytonematode problems of wheat in Hisar and Fatehabad districts of Haryana - A case study --S.N. Nandal and I.J. Paruthi	-- 122
Effect of chemicals on root-knot nematode, <i>Meloidogyne incognita</i> infecting tomato under polyhouse cultivation --H.K. Sharma, Pankaj and Jagan Lal	-- 124
Book review	-- 126